

**PUBLICAÇÃO DE ARTIGOS CIENTÍFICOS COM PARTICIPAÇÃO DA
FUNDAÇÃO HEMOMINAS - ANO DE 2026**

Total de artigos publicados em 2026: 2

HEMOGLOBINOPATIAS (2 artigos)

1- Hematology, Transfusion and Cell Therapy, 48(2):106230, 2026.

doi: <https://doi.org/10.1016/j.htct.2025.106230>

Functional capacity in sickle cell disease: A pilot study with 1-minute sit-to-stand test

Michele Barroso Thomaz¹, Lucas Fernandes Suassuna², Júlia Campos Fabri², Isabela de Oliveira Araújo², Júlia Carneiro Almeida¹, Daniela de Oliveira Werneck Rodrigues¹

1. Fundação Hemominas, Juiz de Fora, MG, Brazil; 2. Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora, MG, Brazil.

Background: Sickle cell disease, the most prevalent monogenic recessive genetic disorder in the world, is characterized by two main pathogenic mechanisms: vaso-occlusion and hemolysis. These characteristics lead to reduced tolerance to physical exertion and, consequently, a reduced functional capacity which can be assessed using the one-minute sit-to-stand test. Complications from sickle cell disease result in poor quality of life, increased absenteeism from school and work, and impaired social interaction.

Method: Between January 2023 and April 2024, a pilot cross-sectional study was conducted with sickle cell disease patients aged from 18 to 60 years. The one-minute sit-to-stand test, Borg's perceived exertion scale, and the SF-36 quality of life questionnaire were utilized. Patients were monitored during the test. The sample was dichotomized based on test performance and SF-36 scores. Furthermore, clinical and demographic variables were analyzed.

Main results: Fifty-eight individuals participated in the final analysis. The mean age was 29.84 ± 11.20 years; 55.1 % were men, and 79.3 % identified themselves as Black or mixed race. The most prevalent genotype was hemoglobin SS (67.2 %), and 77.5 % were taking Hydroxyurea. The group with a better performance in the one-minute sit-to-stand test showed better quality of life as assessed using the SF-36 questionnaire.

Conclusion: Functional capacity is a significant factor in the autonomy and quality of life of patients with sickle cell disease. The one-minute sit-to-stand test is a low-cost and easily applicable test, which can contribute to the assessment of functional capacity in the routine follow-up of these patients.

2- *Annals of Hematology*, 105(4):115, 2026.

doi: <https://doi.org/10.1007/s00277-026-06800-z>

Laboratory and genetic characteristic associated with gallbladder-related outcomes in sickle cell disease in Brazil: results from the REDS-III multicenter cohort study

André Rolim Belisário^{1,2}, Mina Cintho Ozahata³, Isabel Cristina Gomes Moura⁴, Carolina Miranda⁵, Anna Bárbara Carneiro-Proietti⁵, Ester Cerdeira Sabino⁶, Alessandra Ferraz⁷, Cláudia Máximo⁸, Miriam V Flor-Park⁹, Daniela de Oliveira Werneck Rodrigues⁵, Rosimere Afonso Mota⁵, Brian Custer^{10,11}, Shannon Kelly¹², Carla Luana Dinardo¹³; REDS-III Brazil SCD Cohort study and the TOPMed consortium

1. Fundação Hemominas, Belo Horizonte, Minas Gerais, Brazil; 2. Centro de Tecidos Biológicos de Minas Gerais, Lagoa Santa, Fundação Hemominas, Minas Gerais, Brazil; 3. Department of Computer Science - DCC, University of São Paulo, São Paulo, Brazil; 4. Faculdade Ciências Médicas, Minas Gerais, Brazil; 5. Fundação Hemominas, Belo Horizonte, Minas Gerais, Brazil; 6. Faculdade de Medicina (FMUSP), Instituto de Medicina Tropical, Universidade de São Paulo, São Paulo, Brazil; 7. Fundação Hemope, Pernambuco, Brazil; 8. Hemorio, Rio de Janeiro, Brazil; 9. ITACI, Instituto da Criança, HCFMUSP, São Paulo, Brazil; 10. Vitalant Research Institute, San Francisco, CA, USA; 11. Department of Laboratory Medicine, UCSF, San Francisco, CA, USA; 12. UCSF Benioff Children's Hospital, Oakland, CA, USA; 13. Pró-Sangue, São Paulo, Brazil.

Sickle cell disease (SCD) is a hereditary disorder characterized by HBB variants, leading to chronic hemolytic anemia and vaso-occlusion. Hepatobiliary complications, including cholelithiasis, are common but underreported. This study investigated the rates and risk factors for cholelithiasis, cholecystitis, and cholecystectomy in a large Brazilian SCD cohort. Data from 2,778 individuals across six referral centers in the REDS-III Brazilian SCD cohort were analyzed. Clinical, laboratory, and genetic data were obtained retrospectively at enrollment and prospectively during follow-up. Gallbladder-related outcomes were assessed through medical records and imaging. Whole-genome sequencing was performed via the TOPMed program. Genome-wide association analyses used logistic mixed models adjusted for age, sex, genotype, and the first 10 principal components. Cholelithiasis, cholecystitis, and cholecystectomy occurred in 35.9%, 25.1%, and 10.6% of participants, respectively. Indirect bilirubin was consistently associated with all outcomes, while associations with other laboratory variables varied by genotype. Genetic analyses confirmed associations between UGT1A1 variants and bilirubin levels and identified genome-wide associations with cholecystectomy. Novel loci, including FER1L6, LRFN5, and SDK2, were also implicated. These findings indicate a high burden of gallbladder-related disease in Brazilian individuals with SCD and highlight both established and novel genetic pathways that may inform risk stratification and preventive strategies.

COAGULOPATIAS (1 artigo)

1- Haemophilia, 2026 Mar 23. Online ahead of print.

doi: <https://doi.org/10.1111/hae.70244>

Adherence to Clotting Factor Prophylaxis in Adolescent and Adult Males With Haemophilia

Nathália Martins Beserra¹, Ricardo Mesquita Camelo², Alice Oliver Rosa Sacramento³, Eliziane Souza Nascimento⁴, Rosângela Pinheiro Gonçalves Machado⁴, Clarissa Maria Gonçalves Machado⁴, Suzzy Maria Carvalho Dantas⁴, Samuel Gonçalves Machado da Rocha⁴, Luany Elvira Mesquita Carvalho¹⁴, Romélia Pinheiro Gonçalves Lemes⁴

¹ Centro de Hematologia e Hemoterapia do Ceará (HEMOCE), Fortaleza, Brazil ² Faculty of Medicine, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Brazil ³ Fundação Hemominas, Belo Horizonte, Brazil ⁴ Universidade Federal do Ceará (UFC), Fortaleza, Brazil

Background: Clotting factor prophylaxis remains the most prescribed standard-of-care treatment for people with haemophilia (PwH). Prophylaxis prevents bleeds, joint damage, and improves quality of life (QoL), and its success depends on adherence influenced by socioeconomic, behavioural, and disease- or therapy-related factors.

Aim: To identify determinants of clotting factor prophylaxis adherence among PwH followed at a Hemophilia Treatment Center (HTC) in Brazil.

Method: A cross-sectional study was conducted at a Brazilian HTC between June and October 2022. PwH A or B without inhibitors, aged ≥ 14 years and on prophylaxis, were included. Demographic, clinical, and treatment data were collected from records and interviews. QoL was assessed using the SF-36 questionnaire, and adherence using the validated VERITAS-Pro scale.

Main results: Among 78 PwH (median age 30 years), 90% had severe haemophilia and 59% reported joint impairment. Most patients (86%) infused clotting factor three times per week. Median SF-36 score was 70.3, and median VERITAS-Pro score was 44.0, with 90% classified as adherent. Adherence was associated with haemophilia A and infusion frequency >2 times per week, inversely correlated with duration of prophylaxis, and directly associated with better mental health scores. No association was found between adherence and overall QoL.

Conclusion: The main determinants of adherence to clotting factor prophylaxis were having haemophilia A, receiving factor infusions more than twice weekly, shorter duration of prophylaxis, and better mental health.